

Available online at www.sciencedirect.com





Journal of Sport and Health Science xxx (2021) xxx-xxx

Review

Epidemiology of injuries in male and female youth football players: A systematic review and meta-analysis

Francisco Javier Robles-Palazón^a, Alejandro López-Valenciano^{b,*}, Mark De Ste Croix^c, Jon L. Oliver^{d,e}, Alberto García-Gómez^f, Pilar Sainz de Baranda^a, Francisco Ayala^{a,c}

^a Department of Physical Activity and Sport, Faculty of Sport Sciences, Campus of Excellence Mare Nostrum, University of Murcia, Murcia 30720, Spain

^b Centre for Sport Studies, King Juan Carlos University, Madrid 28933, Spain

^c School of Sport and Exercise, University of Gloucestershire, Gloucester GL2 9HW, UK

^d Youth Physical Development Centre, School of Sport and Health Sciences, Cardiff Metropolitan University, Cardiff CF23 6XD, UK

^e Sport Performance Research Institute New Zealand (SPRINZ), Auckland University of Technology, Auckland 0632, New Zealand

^f Operative Research Centre, Miguel Hernández University of Elche, Elche 03202, Spain

Received 4 April 2021; revised 10 June 2021; accepted 21 July 2021

Available online xxx

2095-2546/© 2021 Published by Elsevier B.V. on behalf of Shanghai University of Sport. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/)

Abstract

Background: To conduct a systematic review and meta-analysis of epidemiological data of injuries in male and female youth football players. Methods: Searches were performed in MEDLINE/PubMed, Web of Science, Cochrane Library, and SPORTDiscus databases. Studies were considered if they reported injury incidence rate in male and female youth (< 19 years old) football players. Two reviewers (FJRP and ALV) extracted data and assessed trial quality using the STROBE statement and the Newcastle Ottawa Scale. The Grading of Recommendations Assessment, Development, and Evaluation approach determined the quality of evidence. Studies were combined using a Poisson random effects regression model.

Results: Forty-three studies were included. The overall incidence rate was 5.70 injuries/1000 h in males and 6.77 injuries/1000 h in females. Match injury incidence (14.43 injuries/1000 h in males and 14.97 injuries/1000 h in females) was significantly higher than training injury incidence (2.77 injuries/1000 h in males and 2.62 injuries/1000 h in females). The lower extremity had the highest incidence rate in both sexes. The most common type of injury was muscle/tendon for males and joint/ligament for females. Minimal injuries were the most common in both sexes. The incidence rate of injuries increased with advances in chronological age in males. Elite male players presented higher match injury incidence than sub-elite players. In females, there was a paucity of data for comparison across age groups and levels of play.

Conclusion: The high injury incidence rates and sex differences identified for the most common location and type of injury reinforce the need for implementing different targeted injury-risk mitigation strategies in male and female youth football players.

Keywords: Incidence; Muscle injuries; Severity; Soccer; Young athletes

1. Introduction

Football (soccer) is the most popular sport in the world.¹ Players are required to perform sudden accelerations and decelerations repetitively, rapid changes of directions, and jumping and landing tasks, as well as to be involved in several tackling situations in order to keep possession of or to win the ball.^{2,3} These high-intensity situations, alongside frequent exposure to collisions and contacts, result in a notable increase in injury risk compared to individual sports such as tennis⁴ and gymnastics.⁵ In fact, it has been suggested that football is among the top 5 sports in which players are prone to injury.^{6,7} Injuries are also common events among youth footballers, especially at periods of rapid changes in growth and maturation.^{8–11} Football-related injuries can counter the health-related beneficial effects of sports participation at a young age if a child or adolescent is unable to continue to participate because of the residual effects of injury.¹²

There is a clear necessity to develop and implement measures (e.g., integrative neuromuscular training,13 appropriate rule enforcement, and emphasis on safe play¹⁴) aimed at preventing and reducing the number and severity of football-

*Corresponding author.

Please cite this article as: Francisco Javier Robles-Palazón et al., Epidemiology of injuries in male and female youth football players: A systematic review and meta-analysis, Journal of Sport and Health Science (2021), https://doi.org/10.1016/j.jshs.2021.10.002



E-mail address: alex_tanaco@hotmail.com (A. López-Valenciano).

https://doi.org/10.1016/j.jshs.2021.10.002

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

ARTICLE IN PRESS

related injuries in youth players. However, before implementing any injury prevention measure, it is essential to know the injury profile of youth football.^{15,16} In the past 2 decades, a number of prospective studies have been published describing the incidence and pattern of injuries in youth football players.^{17–27} Recently, a systematic review combined and metaanalysed most of the injury incidences available for elite male youth football and reported overall injury rates of 7.9 and 3.7 time-loss injuries per 1000 h of exposure for players aged under (U) 17 to U21 (i.e., players from around age 17 to age 21) and U9 to U16 (i.e., players from around age 9 to age 16), respectively.²⁸ Furthermore, the same study proposed that a median of 18% (nearly one-fifth) of all reported injuries might be classified as severe (>28 days of absence), with muscle injuries accounting for 37% of all injuries sustained in elite male youth football. However, the systematic review²⁸ also documented a large disparity in injury incidence rates across primary epidemiological studies and pointed out that pooled incidences of injury patterns (i.e., location, type, mechanism, severity of injuries) have not yet been provided for youth foothall

The injury profile in youth male football should not be extrapolated to young female players due to the well-documented anatomical, hormonal, and musculoskeletal sex-related differences.^{29,30} In fact, epidemiological studies have pointed out that male youth footballers seem to be more prone to suffer muscle injuries,^{9,17,18,20–22,24,25,31–34} whereas ligament sprains are the most frequently diagnosed type of injury in female youth footballers.^{27,35} Likewise, disparities in training workloads, medical and performance teams, and physical and mental demands that often exist between elite and sub-elite players and between younger and older age groups might also generate differences in injury incidences according to the level of competition and stages of development.^{3,17,33,36} Indeed. some studies have shown that older adolescent football players who are approaching a professional-league level of play are more susceptible to sustaining injuries than their counterparts playing at a grassroots level.^{37,38}

The potential for differences by sex in youth football-injury profiles requires meta-analytical investigation to identify accurately the most common and severe types of injuries, as well as where (anatomical location) and when (matches or training sessions) they usually occur in these paediatric cohorts. However, to the best of our knowledge, no systematic review and meta-analysis has been published that describes the injury profile of youth football while analysing potential sex differences in injury patterns. Likewise, disparities in training and match demands require the identification of those levels of play and age groups that may present a higher incidence of injury. Therefore, the main purpose of the current study was to conduct a systematic review and meta-analysis quantifying the incidence of injuries in male and female youth football players. The secondary purpose was to determine the overall effects regarding location of injuries, type of injuries, severity of injuries, mechanism of injuries, type of incident, age groups, and level of play.

2. Methods

This systematic review and meta-analysis was carried out following the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) guidelines.³⁹ The PRISMA checklist is presented in Supplementary File 1. The research protocol was registered with PROSPERO (http://www.crd.york.ac.uk/PROSPERO/), registration number CRD42019119279.

2.1. Study selection

Eligibility criteria were established and agreed upon by all authors based on the concept of population, intervention/indicator, comparator/control, outcome, and study design (PICOS)^{39,40} (Supplementary File 2).

Thus, to be included in this systematic review and metaanalysis, studies had to fulfil the following criteria:

- 1) Participants had to be male or female football players \leq 19 years old.
- 2) "Injury" had to be defined in terms of time loss (i.e., an injury that results in a player being unable to take a full part in future football training or match play).^{41,42}
- 3) Eligible studies had to be prospective cohort or randomised control trials (control groups) in order to minimise the occurrence of errors associated with recall.^{41,42} The full text of the article reporting results of the study had to be published in English or Spanish in a peer-reviewed journal before 1 January, 2021.
- 4) Eligible studies had to report either injury incidence rate (IIR) or prevalence among the surveyed players separately by sex or provide sufficient data from which these figures could be calculated through standardised equations.

Studies using injury definitions other than time loss were excluded. Literature reviews, abstracts, editorial commentaries, and letters to the editor were also excluded. Finally, 22 authors were contacted for clarification on raw data extraction^{9,17,24,32,33,35,43-54} and participant information.^{18,19,55,56} Most of the authors contacted (18 out of 22) gave additional details, when requested.^{9,18,19,24,32-34,43,44,47-51,53-56}

2.2. Search strategy

A systematic computerised search was conducted for articles published before 1 January, 2021, in the databases MEDLINE/PubMed, Web of Science, SPORTDiscus, and Cochrane Library. In addition, a complementary search of the reference lists of included articles and a Google Scholar search were also performed. This was done using backward (manually searching the reference list of a journal article) and forward (scanning a list of articles that had cited a given paper since it was published) citation tracking.⁵⁷ When additional studies that met the inclusion criteria were identified, they were included in the final pool of studies. Relevant search terms were used to construct Boolean search strategies, which can be found in Supplementary File 3.

Epidemiology of injuries in youth football

Two authors (FJRP and ALV) used a 2-step process, independently, to select studies for inclusion in the meta-analysis. First, studies were screened based on title and abstract. Second, the full text of the studies identified after the initial screening were reviewed to identify those studies that met the eligibility criteria. A study was excluded immediately when it failed to meet any of the inclusion criteria. Disagreements were resolved through consensus or by consulting a third author (FA).

2.3. Data extraction

A codebook was produced to standardise the coding of each study in order to maximise the objectivity of coding. Each study was codified by 2 different reviewers (FJRP and ALV). The moderator variables of the eligible studies were coded and grouped into 3 categories: (1) general study descriptors; (2) study population; and (3) epidemiological data (injury [including its main characteristics, e.g., location, type, severity, and mechanism, according to Fuller et al.⁴¹] and exposure data). If necessary, the authors of the included studies were contacted to provide clarifications or access to raw data. Operational definitions and moderator variables used in our meta-analysis are shown in Supplementary File 4 and Supplementary File 5, respectively.

The purpose of our meta-analysis was to determine the overall effects, separately, on male and female youth football players of (1) football-related IIR (overall vs. training vs. match); (2) location of injuries (lower extremity vs. trunk vs. upper extremity vs. head and neck); (3) type of injuries (fractures and bone stress vs. joint [non-bone] and ligament vs. muscle and tendon vs. contusions vs. laceration and skin lesion vs. central/peripheral nervous system vs. undefined/other); (4) severity of injuries (slight/minimal [1–3 days] vs. minor/mild [4–7 days] vs. moderate [8–28 days] vs. major/severe [>28 days]); (5) mechanism of injury (overuse vs. traumatic injuries; contact vs. noncontact); (6) new vs. recurrent injuries; (7) age groups (U17–U19, U13–U16, and U12 and below); (8) level of play (sub-elite [low level] vs. elite [high level]); and (9) probabilities of injuries over a season.

With regard to the category level of play, players were classified into 1 of 2 labels: sub-elite or elite. Elite players were defined as follows: football players between 8 and 19 years of age whose performance status was described in the studies as "football academy," "high level," or "elite."^{28,58} Players not described as belonging to a professional youth academy, playing at a high level, or classified as elite were considered to be sub-elite.

The age group category was classified into 3 different labels in order to reflect the taxonomy of children: U12 and below, pubertal adolescents (U13–U16), and post-pubertal adolescents (U17–U19).

2.4. Quality and risk of bias assessment

The reporting quality of included studies was assessed using an adapted version of the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement by von Elm et al.⁵⁹ Supplementary File 6 gives a description of the 22 STROBE criteria designed to assess the quality of the studies included in our meta-analysis. The items and subitems of the STROBE statement were scored as 0 or 1, with a score of 1 given for each checklist item that was properly completed. Using this checklist, a maximum score of 34 would indicate that the article fulfilled requirements for a high-quality publication.

Furthermore, to assess risk of bias of external validity quality, an adapted version of the Newcastle Ottawa Scale (NOS) was used.⁶⁰ This scale contains 8 items and uses a star rating system to indicate the quality of a study (maximum of 8 stars). The higher the number of stars given to an article, the lower the risk of bias. Supplementary File 7 displays a brief description of each item of the adapted version of the NOS used in our meta-analysis.

The data extraction and quality assessments were conducted by 2 reviewers (FJRP, ALV). To assess the inter-coder reliability of the coding process, these 2 reviewers coded 22 (51%) of the included studies randomly (including quality assessment). For the quantitative moderator variables, intra-class correlation coefficients (ICC_{3,1}) were calculated; for the qualitative moderator variables, the Cohen kappa (κ) coefficients were applied. On average, the ICC was 0.84 (range: 0.69–1.00), and the κ coefficient was 0.89 (range: 0.79–1.00), which can be considered highly satisfactory, as proposed by Orwin and Vevea.⁶¹ Inconsistencies between the 2 coders were resolved by consensus, and when they were due to ambiguity in the coding book, they were corrected. As before, any disagreement was resolved by mutual consent in consultation with a third author (FA).

2.5. Quality of the evidence

The quality of the evidence for the overall, training, and match IIRs in male and female youth football players was graded as high, moderate, low, or very low, using a modified GRADE approach. Of the 5 GRADE factors, 4 were used in our meta-analysis: risk of bias, inconsistency, imprecision, and indirectness. The fifth factor, publication bias, is difficult to assess in observational studies due to a lack of registries for these types of studies.⁶² Therefore, we did not take this factor into account in our meta-analysis. The starting point is always the assumption that the pooled or overall result is of high quality. The quality of evidence was subsequently downgraded by 1 or 2 levels per factor to moderate, low, or very low when there was a risk of bias, inconsistency, imprecision, or indirect results.⁶³

2.6. Statistical analysis

IIRs per 1000 hours of player exposure were extracted from the included studies. If IIRs were not reported specifically, they were, if possible, calculated from the available raw data using the following formula:

IIR = $1000 \times (\sum injuries / \sum exposure hours)$

Similar to previous meta-analyses of epidemiology of injuries in sports, ^{60,64,65} data were modelled by a random effects

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

Poisson regression model, as previously described.⁶⁶ The response variable in each meta-analysis was the number of observed injuries, offset by the log of the number of exposure hours (IIR). A random effects term was included to account for the correlation arising from using multiple rows of data from the same study. Factors of interest were included as random effects. The following weighting factor was used: study exposure time (h)/mean study exposure time (h). For the IIR, the overall estimated means for each random effect factor were obtained from the model and then back-transformed to give the IIR, along with 95% confidence intervals (95%CIs that showed negative values were adjusted to 0 for better interpretability). Heterogeneity was evaluated using the I^2 statistic, which represents the percentage of total variation across all studies due to between-study heterogeneity.⁶⁷ The possible influence of the following variables on the model was analysed independently through univariate and multivariate analyses: registration period, year of publication of the study, age of the players, STROBE score, NOS stars, and number of teams included in the study. Sub-analyses separately by sex were carried out when there were at least 3 IIRs (cohorts) coming from a minimum of 2 different studies and the sum of the number of participants involved was more than 30 players.

Where match IIRs were given per 1000 h, post hoc probabilities of injury over a season were determined using the equation developed by Parekh et al.⁶⁸ The Poisson distribution for injury probability has been employed previously in football²⁸ and rugby⁶⁹ studies and describes the frequency of injuries occurring, assuming these injuries occur independently and take place over time or space.⁷⁰ Probability calculations were based on match durations' being between 40 and 90 min, a conservative 30 matches per season, and injuries being independent events.²⁸ Injury probability was calculated separately for male and female players and by age group.

All statistical analyses were performed using the statistical software package R version 2.4.1 (The R Foundation for Statistical Computing, Vienna, Austria) and the "metafor" package.⁷¹

3. Results

3.1. Descriptive characteristics of the studies

A total of 2150 references were identified by all search strategies, of which 43 met the inclusion criteria (resulting in 111 cohort groups, because 19 studies had more than 1 group) (Fig. 1).^{9,10,17–27,31–35,43–51,53–56,72–83} These 43 studies were carried out between 1985 and 2020 and comprised male^{9,10,17,18,20–25,31–34,43–51,54–56,72–80,82,83} and female^{19,26,27,35,43,44,49,51,53,72–74,80,81} players from various (predominantly European) countries. Supplementary File 8 provides a descriptive summary of the characteristics of the included studies.

With regard to the reporting quality of the studies, the mean score obtained with the STROBE quality scale was 23 (minimum = 11; maximum = 32). Regarding the NOS scale, the mean score obtained was 6.5 (minimum = 5; maximum = 8). The quality of evidence according to GRADE was downgraded to moderate (risk of bias and inconsistency) or low (risk of bias, inconsistency, indirectness, and imprecision) for overall, training, and match IIR outcomes in males and females, respectively. The detailed data for STROBE, NOS, and GRADE scales are presented in Supplementary File 9, Supplementary File 10, and Supplementary File 11, respectively.

3.2. Meta-analyses

In the various meta-analyses we carried out, the effect sizes exhibited a moderate to large heterogeneity (based on the Q statistics and the I^2 indices), supporting the decision to apply random-effects models.

None of the following had an impact on IIRs: the registration period (i.e., the period of time/year when the data collection process was carried out), the year of publication of the study, age, STROBE score, NOS stars, or number of teams. Hence, the subsequent sub-analyses were not adjusted to these variables.

3.2.1. Injury incidence: overall, training and match

Males. Thirty-three studies (38 cohorts) reporting overall IIRs, 9,10,17,18,20-25,31-34,43,45-51,54-56,73,75-79,82,83 25 studies cohorts) reporting (30 training IIRs,^{18,20–22,24,25,31–34,43,45,47–51,54,56,75,77–79,82,83} and 29 studies (34 cohorts) reporting match IIRs^{18,20-22,24,25,31-34,43-45,47-51,54,56,72,74,75,77-80,82,83} in male youth football players were included in our meta-analysis, comprising a total of 7495 injuries and about 25,600 different players. The random effect models showed an overall IIR of 5.70 injuries/1000 h (95%CI: 4.54–6.86, $I^2 = 98.0\%$; quality of evidence = moderate), a training IIR of 2.77 injuries/1000 h (95%CI: 2.04–3.50, $I^2 = 97.0\%$; quality of evidence = moderate), and a match IIR of 14.43 injuries/1000 h (95%CI: 11.00-17.85, $I^2 = 97.0\%$; quality of evidence = moderate). Fig. 2 and Fig. 3 display the forest plots with the training and match IIRs, respectively, for males in the analysed studies.

Females. Nine studies (11 cohorts) reporting overall IIRs, ^{26,27,35,43,49,51,53,73,81} 5 studies (7 cohorts) reporting training IIR, ^{27,43,49,51,53} and 10 studies (12 cohorts) reporting match IIRss^{19,27,43,44,49,51,53,72,74,80} in female youth football players were included in our meta-analysis, comprising a total of 2179 injuries and about 9600 different players. The random effect models showed an overall IIR of 6.77 injuries/1000 h (95%CI: 5.01-8.52, $I^2 = 94\%$; quality of evidence = low), a training IIR of 2.62 injuries/1000 h (95%CI: 1.16-4.08, $I^2 = 90\%$; quality of evidence = low). Fig. 4 and Fig. 5 display the forest plots with the training and match IIRs, respectively, for females in the analysed studies.

3.2.2. Location of injury

Males. Twenty-four studies reported injury location and lower-extremities-region categories (based on Fuller et al.⁴¹) in males. $^{9,17,18,20,21,23-25,31-34,43,45,47,48,50,54,56,77-79,82,83}$

Epidemiology of injuries in youth football



Fig. 1. Flow chart of the selection of studies for the meta-analysis. ^aNo injury definition (n = 2), full-text not available (n = 2) and incidence for football players reported jointly with other sports (n = 1).

Lower-extremity iniuries had the highest IIR (IIR = 4.08 injuries/1000 h; 95%CI: 2.92-5.24, $I^2 = 99.5\%$) compared to injuries in other body regions. The upper limbs region was the second most commonly injured region (IIR = 0.29 injuries/1000 h; 95%CI: 0.20-0.39, $I^2 = 94.7\%$); the trunk region was third (IIR = 0.25 injuries/1000 h; 95%CI: 0.17-0.34, $I^2 = 92.9\%$), and injuries to the head and neck region had the lowest IIR (IIR = 0.08 injuries/1000 h; 95%CI: 0.04-0.12, $I^2 = 88.5\%$). Regarding lower-extremity injuries, the thigh showed the highest IIR (IIR = 1.21, 95%CI: 0.74 - 1.69, $I^2 = 99.1\%$), followed by ankle ((IIR = 0.91, 95%CI: 0.64–1.18, $I^2 = 97.6\%$), knee (IIR = 0.75, 95%CI: 0.53-0.97, $I^2 = 96.6\%$), hip/groin (IIR = 0.73, 95%CI: $0.45 - 1.00, I^2 = 98.1\%$), lower leg/Achilles tendon (IIR = 0.37, 95%CI: 0.24–0.50, $I^2 = 94.4\%$), and foot/toe (IIR = 0.31, 95%CI: 0.19-0.43, $I^2 = 94.9\%$).

Females. Only 5 studies reported injury location and lowerextremities region categories in female youth footballers.^{26,27,35,43,53} The trend was similar to that for males, with lower extremities having the highest IIR (IIR = 6.54 injuries/1000 h; 95%CI: 4.68-8.40, $I^2 = 91.4\%$), followed by trunk (IIR = 0.68 injuries/1000 h; 95%CI: 0.54-0.82, $I^2 = 0\%$) and upper limbs (IIR = 0.26 injuries/1000 h; 95%CI: 0.12–0.39, $I^2 = 51.0\%$), with the lowest IIR related to head and neck injuries (0.13 injuries/1000 h; 95%CI: 0.00-0.34, $I^2 = 68.2\%$). With regard to lower-extremity injuries, ankle (IIR = 1.52, 95%CI: 1.19–1.86, $I^2 = 64.0\%$) and knee (IIR = 1.49, 95%CI: 0.90–2.08, $I^2 = 89.3\%$) showed the highest IIRs, followed by thigh (IIR = 1.06, 95%CI: 0.55-1.56, $I^2 = 91.0\%$), lower leg/Achilles tendon (IIR = 0.68, 95%CI: 0.28–1.08, I^2 = 90.2%), hip/groin (IIR = 0.57, 95%CI: 0.19-0.96, $I^2 = 91.9\%$), and foot/toe (IIR = 0.40, 95%CI: $0.31 - 0.49, I^2 = 0\%$ (Fig. 6).

3.2.3. Type of injury

Males. Fifteen studies reported the type of injury among male players.^{9,17,18,21,24,25,31–34,43,47,48,50,82} The most common type of injury grouping was muscle/tendon (IIR = 1.92 injuries/1000

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637

638

639

640

641

642 643

644

645

646

647

648

649

650

651

652

653

654

655

656

657

658

660

661

662

663

664

665

666

667

668

669

670

671

672

673

674

50



Fig. 2. Training injury incidence in male youth football players, with 95% confidence intervals.

Study 95%CI 3.60 (2.86 - 4.34) Andreasen et al. (1992)72 Azuma et al. (2020)47 11.90 (8.43 - 15.38) Bianco et al. (2016)24 2.84(1.65 - 4.03)6.80(3.58 - 10.02)Brito et al. (2011)34 Brito et al. (2012)33 4.70(3.48 - 5.92)Cezarino et al. (2020)22 6.41 (4.16 - 8.67) Chena-Sinovas et al. (2019)21 8.88 (6.97 - 10.79) Ergün et al. (2013)31 30.40 (15.01 - 45.79) Frisch et al. (2011)32 23.50 (18.15 - 28.85) Hägglund et al. (2009a)51 16.30(5.00 - 27.60)Hägglund et al. (2009b)51 27.80 (13.24 - 42.36) Hägglund et al. (2009c)51 21.83(8.93 - 34.72)Hägglund et al. (2009d)51 20.70 (7.86 - 33.54) Hägglund et al. (2009e)51 12.58 (2.51 - 22.64) Hägglund et al. (2009f)51 28.60 (13.06 - 44.14) 37.20 (30.22 - 44.18) Hawkins et al. (1999)56 Imai et al. (2018)54 6.24 (2.55 - 9.93) Junge et al. (2013)74 28.39 (24.93 - 31.84) Junge et al. (2004)75 16.20 (12.06 - 20.34) Kakavelakis et al. (2003)77 5.60(4.50 - 6.70)Kuzuhara et al. (2017)43 5.44 (2.48 - 8.39) Nielsen et al. (1989)78 14.40 (6.92 - 21.88) Nilsson et al. (2016)20 15.50 (8.34 - 22.66) 14.22 (11.74 – 16.70) Nogueira et al. (2017)25 Owoeye et al. (2014)79 20.30(15.64 - 24.96)Raya-González et al. (2019)18 8.84 (7.22 - 10.45) 9.72 (8.21 - 11.22) Rommers et al. (2020)44 Schmidt-Olsen et al. (1985)80 7.40(5.99 - 8.81)Sieland et al. (2020)45 5.00 (3.27 - 6.73) Soligard et al. (2012)44 3.87 (3.29-4.45) Sprouse et al. (2020)49 26.53 (22.71 - 30.34) Waldén et al. (2007)82 30.40 (15.04 - 45.76) Wik et al. (2020)50 31.98 (29.10 - 34.87) Zarei et al. (2020)83 10.01(6.00 - 14.02)14.43 (11.00 - 17.85) Total Heterogeneity: $\chi^2_{\rm 33}$ = 1055.23 ($p < 0.001), \ l^2$ = 97% 0 10 20 30 40

Fig. 3. Match injury incidence in male youth football players, with 95% confidence intervals.

6

608

609

610

611

612

613

614

615

616

617

Please cite this article as: Francisco Javier Robles-Palazón et al., Epidemiology of injuries in male and female youth football players: A systematic review and meta-analysis, Journal of Sport and Health Science (2021), https://doi.org/10.1016/j.jshs.2021.10.002

F.J. Robles-Palazón et al.

Epidemiology of injuries in youth football



Fig. 4. Training injury incidence in female youth football players, with 95% confidence intervals.



Fig. 5. Match injury incidence in female youth football players, with 95% confidence intervals.

h; 95%CI: 1.26–2.58, $I^2 = 99.0\%$), followed by joint (non-bone) and ligament (IIR = 0.97 injuries/1000 h; 95%CI: 0.64-1.30, $I^2 = 97.4\%$) and contusions (IIR = 0.84 injuries/1000 h; 95%CI: 0.37 - 1.30. $I^2 = 99.3\%$). Fracture and bone stress (IIR = 0.43 injuries/1000 h, 95%CI: 0.02-0.84, $I^2 = 99.7\%$), undefined/other (IIR = 0.27 injuries/1000h; 95%CI: 0.04-0.50, $I^2 = 99.5\%$), central/peripheral nervous system (IIR = 0.06 injuries/1000 h; 95%CI: 0.00-0.12, $I^2 = 95.6\%$) and laceration and skin lesions (IIR = 0.03 injuries/1000 h; 95%CI: 0.01-0.05, $I^2 = 66.0\%$) were the least common types of injury.



Fig. 6. Location of injuries in male (left side) and female (right side) youth football players. The upper boxes (with solid lines extending from them) represent the incidence of injury for main groups (head and neck, upper limbs, trunk, lower limbs), whereas the lower boxes (with dashed lines extending from them) represent the incidence of injury for lower extremities categories (hip/groin, thigh, knee, lower leg/Achilles tendon, ankle, foot/toe).

Females. Only 3 studies reported type of injury in females players and were the only 3 pooled in our metaanalysis.^{27,35,43} Unlike injuries in males, joint (non-bone) and ligament injuries (IIR = 2.36 injuries/1000 h; 95%CI: 1.62-3.11, $I^2 = 59.0\%$) were the most common type of injury among females, followed by muscle and tendon injuries (IIR = 2.01 injuries/1000 h; 95%CI: 1.75-2.28, $I^2 = 0\%$), contusions (IIR = 0.93 injuries/1000 h; 95%CI: 0.61-1.25, $I^2 = 44.6\%$), undefined/other (IIR = 0.84 injuries/1000 h; 95%CI = 0.49-1.20, $I^2 = 57.0\%$), and fracture and bone stress injuries (IIR = 0.27 injuries/1000 h; 95%CI: 0.17-0.36, $I^2 = 0\%$). No laceration and skin lesions or central/peripheral nervous system injuries were registered.

3.2.4. Severity of injury

Males. Twenty-one studies (26 cohorts) reported severity of injury in males.^{9,17,18,20,21,23,25,31–34,45–48,50,51,56,79,82,83} Minimal injuries (IIR = 1.88 injuries/1000 h; 95%CI: 1.13–2.64, $I^2 = 99.8\%$) were the most common, followed by moderate (IIR = 1.74 injuries/1000 h; 95%CI = 1.30–2.18, $I^2 = 98.0\%$), mild (IIR = 1.13 injuries/1000 h; 95%CI: 0.77–1.49, $I^2 = 98.5\%$), and severe (IIR = 0.78 injuries/1000 h; 95%CI: 0.56–1.00, $I^2 = 96.4\%$). Additionally, a total of 11 studies^{9,18,23–25,31–34,46,83} reported an average of 15.5 days lost per injury by male footballers, with an overall injury burden of 96.47 injury days/1000 h of football exposure (95%CI: 49.86–143.08, $I^2 = 100\%$).

Females.	Only 3	studies (5 cohorts)	reported on the	severity of
injuries	in	females. ^{19,51,53}	Minimal	injuries

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826

827

828

829

830

831

832

833

834

835

836

837

838

839

840

841

842

843

844

845

ARTICLE IN PRESS

(IIR = 3.60 injuries/1000 h; 95%CI: 0.68–6.53, I^2 = 82.3%) were also the most usual in females, followed by moderate (IIR = 1.52 injuries/1000 h; 95%CI: 1.18–1.87, I^2 = 0%), severe (IIR = 1.25 injuries/1000 h; 95%CI: 0.57–1.94, I^2 = 43.1%), and mild (IIR = 0.76 injuries/1000 h; 95%CI: 0.51–1.00, I^2 = 0%). The paucity of data prevented the calculation of pooled estimates for the injury burden among female footballers.

3.2.5. Mechanism of injury

Males. In our meta-analysis, 16 studies (19 cohorts) provided data for the comparison of overuse injuries to traumatic (acute) injuries among males.^{9,18,20,21,23,25,31–33,49–51,54,56,79,82} The IIR for traumatic injuries (IIR = 5.50 injuries/1000 h, 95%CI: 3.98-7.01) was higher than for overuse injuries (IIR = 1.10 injuries/1000 h, 95%CI: 0.68-1.53). In relation to the mechanism of injury, 15 studies (18 cohorts) reported data that allowed us to compare contact injuries to noncontact injuries among males. ^{9,21,25,31–33,45,47,49–51,55,56,79,82} Males showed a slightly higher IIR for noncontact injuries (IIR = 3.48 injuries/1000 h, 95%CI: 2.35-4.62) than for contact injuries (IIR = 2.77 injuries/1000 h, 95%CI: 1.93-3.61).

Females. Eight studies (9 cohorts) provided data for the comparison of overuse injuries to traumatic (acute) injuries among females.^{19,26,27,35,49,51,53,81} Similar to the IIR for males, the IIR for traumatic injuries (IIR = 4.55 injuries/1000 h, 95%CI: 3.72-5.38) among females was higher than for overuse injuries (IIR = 1.56 injuries/1000 h, 95%CI: 0.80-2.31). Four studies (5 cohorts) reported data that allowed us to compare contact noncontact injuries to injuries among females.49,51,53,81 Similar IIRs for noncontact injuries (IIR = 2.39 injuries/1000 h, 95%CI: 1.80-2.98) and for contact injuries (IIR = 1.92 injuries/1000 h, 95%CI: 1.67-2.17) were found.

3.2.6. New vs. recurrent injuries

Males. Included in our analysis were 11 studies (14 cohorts) that aimed to compare the IIR for new injuries to the IIR for recurrent injuries among males.^{9,24,25,31–34,48,50,51,56} The IIR for new injuries (IIR = 5.87 injuries/1000 h, 95%CI: 3.89-7.84) was higher than that for recurrent injuries (IIR = 0.81 injuries/1000 h, 95%CI: 0.38-1.25).

Females. Five studies (6 cohorts) provided data that allowed us to compare the IIR for new injuries to the IIR for recurrent injuries among females.^{26,27,35,51,81} Similar to the IIR for males, the IIR for new injuries (IIR = 5.12 injuries/1000 h, 95%CI: 3.64–6.61) was higher than the IIR for recurrent injuries (IIR = 1.42 injuries/1000 h, 95%CI: 0.29–2.55) among female footballers.

3.2.7. Age groups

Males. Concerning the age of football players, we categorised footballers into 3 groups: U12 and below, U13–U16, and

U17-U19. For 20 studies (58 males, cohorts)^{9,18,21–25,31,33,43,46,49–51,54–56,77,78,82} provided data that allowed us to compare the overall IIR; 16 studies (46 cohorts) $^{9,18,21,22,24,25,31,43,49-51,54,56,77,78,82}$ provided data that allowed us to compare training IIRs; and 19 studies (55 cohorts)^{9,18,21,22,24,25,31,43,44,49-51,54,56,74,77,78,80,82} provided data that allowed us to compare match IIRs. The U17-U19 male age group showed the highest overall IIR (IIR = 7.54 injuries/1000 h; 95%CI: 5.62-9.47, $I^2 = 97\%$), followed by the U13-U16 male group (5.35 injuries/1000h, 95%CI: 3.73-6.98, $I^2 = 98\%$), and the U12 male group (IIR = 1.61 injuries/1000 h; 95%CI: 0.76–2.45, $I^2 = 85\%$). In particular, the mean IIRs in training decreased from the U17-U19 age group (IIR = 3.51 injuries/1000 h; 95%CI: 2.15-4.87, $I^2 = 91\%$) to the U13-U16 age group (IIR = 3.39 injuries/1000h, 95%CI: 2.20–4.57, $I^2 = 95\%$) to the U12 age group (IIR = 1.07 injuries/1000 h; 95%CI: 0.40–1.74, $I^2 = 72\%$). The match IIRs for each age group were. in descending order: U17-U19 (IIR = 20.05 injuries/1000 h; 95%CI: 15.48–24.62, $I^2 = 93\%$); U13-U16 (IIR = 13.67 injuries/1000 h; 95%CI: 8.49-18.86, $I^2 = 95\%$; and U12 (IIR = 2.60 injuries/1000 h; 95%CI: $0.60 - 4.59, I^2 = 77\%$).

Females. Only 2 studies (5 cohorts) provided data that allowed us to compare overall and training IIRs for females,^{49,51} and only 6 studies (15 cohorts) provided data that allowed us to compare match IIRs.^{19,44,49,51,74,80} The U17–U19 female age group had an overall IIR of 6.25 injuries/1000 h (95%CI: 4.68–7.81, $I^2 = 38\%$), a training IIR of 3.08 injuries/1000 h (95%CI: 2.18–3.98, $I^2 = 40\%$), and a match IIR of 20.94 injuries/1000 h (95%CI: 14.27–27.62, $I^2 = 78\%$). The U13–U16 female age group reported a match IIR of 12.67 injuries/1000 h (95%CI: 5.41–19.94, $I^2 = 89\%$). The scarcity of studies reporting overall, training and match injury IIRs for the U12 and below female age group, and overall and training IIRs for the U13–U16 female age group, prevented further sub-analyses for these groups.

3.2.8. Level of play

Males. Regarding the level of play, studies were classified into 2 groups: sub-elite and elite. Ten studies reported overall IIRs, ^{21,25,32–34,43,47,54,75} 9 studies reported training IIRs, ^{21,25,32–34,43,47,54,75} and 9 studies reported match IIRs^{21,25,32–34,43,47,54,75} in sub-elite players. The random effect models showed an overall IIR of 4.77 injuries/1000 h (95%CI: 2.63–6.90, $I^2 = 98\%$), a training IIR of 2.83 injuries/1000 h (95%CI: 1.39–4.27, $I^2 = 96\%$), and a match IIR of 10.63 injuries/1000 h (95%CI = 5.98–15.28, $I^2 = 93\%$).

For its part, the elite level was represented by 20 (25 cohorts) overall IIR studies, 9,10,17,18,20,22–24,31,45,46,48–51,55,56,79,82,83 14 (19 cohorts) training IIR studies,^{18,20,22,24,31,45,48-51,56,79,82,83} and 16 studies (21 cohorts) from competitions.^{18,20,22,24,31,44,45,48–51,56,74,79,82,83} The random effect models showed an overall IIR of 6.19 injuries/1000 h (95%CI: 4.56-7.82, $I^2 = 99\%$), a training IIR of

901

Epidemiology of injuries in youth football

2.68 injuries/1000 h (95%CI: 1.64–3.72, $I^2 = 98.0\%$), and a match IIR of 17.91 injuries/1000 h (95%CI: 12.99–22.83, $I^2 = 98\%$).

Females. Three studies (4 cohorts) reported overall IIR^{26,35,43} in sub-elite female players, with the random effect models displaying a total of 7.86 injuries/1000 h of football exposure (95%CI: 3.27-12.44, $I^2 = 78\%$). An insufficient number of studies were found to estimate training and match IIRs in sub-elite female players.

On the other hand, 4 studies (6 cohorts) reported overall, 26,27,49,51 3 studies (4 cohorts) reported training, 27,49,51 and 5 studies (6 cohorts) presented match 27,44,49,51,74 IIRs in elite female players. The overall IIR was 6.49 injuries/1000 h (95%CI: 5.76-7.23, $I^2 = 50\%$), 3.24 injuries/1000 h of training (95%CI: 1.60-4.89, $I^2 = 79\%$), and 18.13 injuries/1000 h of match (95%CI: 9.43-26.82, $I^2 = 98\%$).

3.2.9. Probability of Injury

The overall injury probability during 1 season was 47% and 43% for male and female youth players, respectively. Independent of sex, the highest injury probability was found for the U17–U19 age groups (56% in males and 58% in females) and was lowest for the U12 age group (7% in males and 18% in females). The U13-U16 age group had an injury probability of 39% for males and 30% for females. Supplementary File 12 provides a descriptive summary of the probabilities of injury for both male and female cohorts.

4. Discussion

Both the methodology and the statistical analyses used in our study were identical to those used in the systematic reviews and meta-analyses conducted by Lopez-Valenciano et al.^{60,84} for adult men (elite football players) and women (subelite and elite football players). Although injury profile comparisons between youth and adult football players are possible, our comparisons should be interpreted with a certain degree of caution due to inter-meta-analyses differences in the number of cohorts and quality of the studies included in each analysis.

4.1. Injury incidence: Overall, training, and match

The main findings in our study indicate that the overall, training, and match IIRs in male youth football players (5.7 injuries/1000 h, 2.8 injuries/1000 h, and 14.4 injuries/1000 h of overall, training and match exposure, respectively) and female youth football players (6.8 injuries/1000 h, 2.6 injuries/1000 h, and 15.0 injuries/1000 h of overall, training and match exposure, respectively) are higher than the IIRs found in previous studies related to other youth team sports such as handball (2.9 injuries/1000 h, 0.9 injuries/1000 h, and 9.9 injuries/1000 h of overall, training and match exposure, respectively)⁸⁵; basketball (1.3 injuries/1000 h, 0.5 injuries/1000 h, 11.2 injuries/1000 h of overall, training and match exposure, respectively)⁸⁶; and volleyball (2.4 injuries/1000 h of match exposure).⁸⁷ Furthermore, the probability of youth football players' sustaining a time-loss injury during a season was 47%

for male players and 43% for female players. These probability-of-injury scores are higher than the 28% score reported for child and adolescent rugby players during a rugby season.⁶⁹ The high IIRs and probability scores found for youth footballers in our meta-analysis reinforce the need for implementing targeted injury risk mitigation strategies in youth football.

For adult football players^{60,84} and players in other youth team sports (independent of the players' sex), such as handball,⁸⁵ basketball,⁸⁶ volleyball,⁸⁷ and rugby,⁶⁹ match IIRs have always been found to be significantly higher than training IIRs. A number of studies have attributed the difference in IIRs between match and training to several factors, including the higher physical playing demands during matches compared to the demands of training sessions, the greater variability and uncertainty in game demands when competing against rivals (compared to the familiarity of training with teammates), the number of contacts and collisions during matches, and the fatigue generated during the course of a match.^{38,88,89}

4.2. Location and type of injuries

Similar to the rates that have been reported for adult footballers, lower-extremity injuries had the highest IIRs compared to the other body regions among both male and female youth football players (4.1 injuries/1000 h and 6.5 injuries/1000 h, respectively).

The locations of the most frequently reported injuries in male and female youth footballers were slightly different. In male players, the thigh (1.2 injuries/1000 h) and ankle (0.9 injuries/1000 h) were the anatomical regions where injuries occurred most frequently, whereas the knee (1.5 injuries/1000 h) and ankle (1.5 injuries/1000 h) were the regions where injuries were reported most frequently among females. The higher knee and ankle IIRs found for female youth football players in our meta-analysis may be explained by the fact that females sustained twice as many joint (non-bone) and ligament injuries as their male counterparts (2.4 injuries/1000 h for females and 1.0 injuries/1000 h for males). This higher susceptibility for sustaining joint and ligament injuries observed in female youth football players in comparison with their male counterparts has also been found in adult football players. Sexrelated differences in core and lower extremity neuromuscular control, joint laxity, hormonal regulation, biomechanics, and anatomy^{29,30,90} have been suggested (among other factors) as reasons why female athletes are more prone to suffering joint (non-bone) and ligament injuries, mainly around the knee and ankle joints. Because of the lack of epidemiological studies reporting IIRs separately for joints (non-bone) and ligaments (e.g., anterior cruciate ligament of the knee and anterior inferior tibiofibular ligament of the ankle) among youth footballers, a subanalysis aimed at identifying the most commonly injured joint (non-bone) and ligament was not possible. However, previous studies have consistently reported that ankle sprains were the most frequent joint and ligament injuries diagnosed in youth football players, independent of the sex of the players.^{8,18,27,35}

1017

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

1029

1030

1031

1032

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

In our meta-analysis, the thigh was the area most frequently injured in male football players. However, no sex-related differences were found in the magnitude of thigh IIRs $(\sim 1.1 \text{ injuries}/1000 \text{ h for both male and female players})$. This circumstance strongly correlates with the fact that both male and female youth football players also presented analogous muscle IIRs (~ 2 injuries/1000 h). The link between these 2 IIRs can be found in the fact that hamstring and quadriceps muscle injuries, both operationally located in the thigh,⁴¹ have been consistently reported as the most frequently diagnosed injuries in youth football players (and also in adult players).^{8,38,91} However, the very limited number of studies available that reported IIRs separately by muscle group prevented us from calculating pooled estimates for hamstring and quadriceps muscle injuries. In contrast, it should be noted that, among adult football players, men and women did not report similar muscle injury rates. In particular, male footballers presented muscle IIRs that were twice as high as women's IIRs (4.6 injuries/1000 h vs. 1.8 injuries/1000 h, respectively), which might be attributed to the larger intersex differences in physical match demands (e.g., number of high-intensity actions performed) that are evident in elite football.⁹²

Interestingly, the IIRs related to trunk injuries were more than twice as high for female footballers as for male footballers (0.7 injuries/1000 h *vs*. 0.3 injuries/1000 h, respectively) but were still relatively low for both sexes. A more erect posture during landing has been evidenced in females, which could overload not only the lower limbs but also the trunk area.⁹⁰ Consequently, this may increase the risk of trunk injuries (e.g., spondylolisthesis) for females. Therefore, it would be advisable for prevention programs for females to focus on core strength also.

Current international research has also given particular attention to head injuries that involve the nervous system (i.e., concussions and traumatic brain injuries). Our results in this area showed the lowest IIRs for head and neck injuries (0.1 injuries/1000 h) and for injuries to the central/peripheral nervous system (< 0.1 injuries/1000 h) in both males and females, which matches the findings of previous large-scale investigations.93,94 However, these injuries might be underdiagnosed frequently due to inconsistencies in the interpretation and reporting of the symptoms.⁹⁵ Thus, the use of a definition of a time-loss injury may have reduced the proportion of concussion injuries pooled in our research. Future prospective studies using a more accurate injury definition, as well as a recognition of and reporting on this type of injury, are needed to analyse the evidence on the incidence of concussions among youth footballers.

4.3. Severity and mechanisms of injuries

Although injuries occur frequently in youth football players, the majority of injuries, fortunately, appear to be of minimal severity (1-3 days lost). However, it should be highlighted that the IIRs for moderate injuries (1.7 injuries/1000 h for males and 1.5 injuries/1000 h for females) and for severe injuries (0.8 injuries/1000 h for males and 1.3 injuries/1000 h for females) found

F.J. Robles-Palazón et al.

in our meta-analysis may be considered problematic. In practical terms, our findings might imply that in a typical youth football squad comprising 20 players, a coach could expect 2 highburdensome injuries (> 28 days of time loss) per season (value calculated using the data provided in original studies $^{9,18,20,21,23,25,31-34,45,47,48,50,51,53,56,79,82,83}$). Results of our study have revealed that a great proportion of injuries in male and female youth footballers might have traumatic and noncontact mechanisms and, as such, they can be regarded as preventable. The implementation of comprehensive injury prevention programs aimed at improving movement competency and physical fitness among youth footballers has demonstrated that this can be a successful approach to reducing the number of moderate and severe noncontact injuries in children and adolescents.^{53,96} Previous studies have demonstrated that 10-15 min of neuromuscular training activities 2 to 3 times weekly reduces noncontact injuries by 45% in youth football players.97

Although injuries to adult football players can have negative effects on a team and its success rate,98 the impact of injuries on the development of youth football players has yet to be established. However, it may be assumed that at young ages, being away from football play for more than 28 days may not only negatively influence the short-term tactical, technical, and physical performance of youth football players but may also impair their long-term development, health outcomes, and future career opportunities.^{12,99} Because the studies included in our meta-analysis reported only IIRs, not the average number of days lost from football (time loss) by location and type of injury, it was not possible for us to calculate the injury burden and, thus, build a risk matrix. A risk matrix would have helped to identify the importance (i.e., burden) of each football-related injury and provided information that could have helped prioritise injury-prevention measures used in applied football environments. However, based on the findings from previous studies,^{46,100} the most burdensome injuries in youth football may well be quadriceps and hamstring muscle injuries, knee ligament injuries (anterior cruciate ligament tears) and growth-related injuries (Osgood-Schlatter and Sinding-Larsen diseases). In terms of the severity and mechanisms of injuries described for youth football players, this injury pattern is very similar to that reported by López-Valenciano et al.^{60,84} for adult footballers.

4.4. New vs. recurrent injuries

As expected, and similar to the findings reported elsewhere concerning adult football players,^{60,84} the IIRs for recurrent injuries in youth footballers is lower than the IIRs for new injuries (0.8 injuries/1000 h for males and 1.4 injuries/1000 h for females *vs.* 5.9 injuries/1000 h for males and 5.1 injuries/1000 h for females, respectively). Likewise, there were no sexrelated differences in new and recurrent injuries in either youth or adult football players. However, it should be highlighted that the ratio of new injuries to recurrent injuries was higher among youth players (7.4 for male youths *vs.* 5.4 for male adults⁶⁰ and 3.6 for female youths *vs.* 2.6 for female adults⁸⁴).

1128

1129

1130

Please cite this article as: Francisco Javier Robles-Palazón et al., Epidemiology of injuries in male and female youth football players: A systematic review and meta-analysis, Journal of Sport and Health Science (2021), https://doi.org/10.1016/j.jshs.2021.10.002

Epidemiology of injuries in youth football

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

1164

1165

1166

1167

1168

1169

1170

1171

1172

1173

1174

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

The lower recurrent IIRs in youth players in comparison with their adult counterparts may indicate that at young ages there is not such high pressure to return to play as soon as possible, contributing to improved rehabilitation.^{58,101} On the other hand, having a previous history of injury is 1 of the few evidence-based predictors available in the literature for the most common football-related injuries (i.e., hamstring and knee injuries).¹⁰²⁻¹⁰⁴ As a consequence of having more experience in playing football, adult footballers may present a higher likelihood of having suffered previous injuries than youth players; hence, adults may be at a higher risk of injury recurrence.^{105,106} This circumstance has led some researchers to suggest that the main purpose of injury-risk-mitigation strategies in youth football should be to delay, as much as is possible, the occurrence of the first injury.^{105,107} Longitudinal studies that track IIRs through the academy setting and into professional environments might help to elucidate whether there is a consequence of repeated injuries during growth and maturation.¹⁰⁸

4.5. Age groups

Results from the various age groups, representing differing periods of childhood and adolescence, suggest potential interactions among maturity, sex, training, and competition with IIRs. In males, the overall IIR increased among players who are likely to be prepubertal (U12), circapubertal (U13–U16), or postpubertal (U17–U19),¹⁰⁹ with overall IIRs of 1.6 injuries/1000 h, 5.3 injuries/1000 h, and 7.5 injuries/1000 h of football exposure, respectively. This was driven by a high IIR for matches, which increased by approximately 10 injuries between each consecutive age interval (2.6 injuries/1000 h vs. 13.7 injuries/1000 h vs. 20.0 injuries/1000 h). The changing profile of the IIR is likely to be attributable to both maturation effects and increasing demands of training and competition in older age groups. Young children have an immature neuromuscular and metabolic system, with a lower muscle mass, more compliant muscle-tendon structures, and less ability to recruit fast-twitch fibres, with an underdeveloped anaerobic system and a greater reliance on aerobic metabolism.¹¹⁰ All these factors mean that immature players work less explosively and that they generate and have to tolerate lower levels of force, thus exposing themselves to lower levels of risk. At the same time, they experience lower levels of fatigue during intermittent work and are able to recover from fatigue more quickly.¹¹¹ This is reflected in the fact that U12 players have low overall and low match IIRs. Adolescent players experience a period of rapid physical development that will result in gains in both size and fitness, but this developmental period can be accompanied with temporarily disrupted motor coordination.¹¹² Consequently, adolescent players may begin to expose themselves to a greater intensity and volume of exercise within training and match play and may display aberrant movement mechanics while also being more susceptible to growth and overuse injuries.^{8,10,113} They may also have a reduced ability to recover between matches.¹¹⁴ All these

factors may contribute to a higher IIR for adolescent players compared to prepubertal players.

Players continue to develop physically into late adolescence and early adulthood and will likely continue to increase their abilities to work at high intensities, completing more accelerations, decelerations, and greater total distances during competition compared to younger players.¹¹⁵ The increased physical demands and longer duration of match play mean that players in the older age groups are exposing themselves to more risk during a game. Simultaneously, players transitioning to older age groups (U17-U19) are likely to experience a great increase in training load as they begin to train on full-time professional contracts,⁸ and these spikes in workload have been suggested to contribute to injuries among youth football players.^{8,116} These increases in IIRs across players' age groups are also evident when compared with the results reported by López-Valenciano et al.⁶⁰ for adult footballers, where IIRs reach up to 8.1 injuries/1000h, 3.7 injuries/1000h and 36.0 injuries/1000h for overall, training and match exposure, respectively.

There was a paucity of data available to compare IIRs across age groups for female players. Only 2 studies reported overall and training IIRs for females in the U17–U19 age groups,^{49,51} and although a few others have presented match injury data for females in the U13–U16 and U17–U19 cohorts, most of these studies correspond to football tournaments,^{19,44,51,74,80} and 1 was published in 1985.⁸⁰ Based on the available information, females who were in the U17–U19 group experienced a higher incidence of match-related injuries than U13–U16 females (20.9 injuries/1000 h *vs.* 12.7 injuries/1000 h, respectively), which is similar to the increase described for males. However, more research with longer follow-up periods is needed to confirm the potential differences between age groups in females, especially across a range of maturational stages.

4.6. Level of play

The findings from our study also indicate that elite (highlevel) male players present a higher match IIR (17.9 injuries/1000 h) than their sub-elite (less skilled) peers (10.6 injuries/1000 h). These observed differences according to the level of play may be partially explained by the fact that elite players perform more high-intensity actions during competitions and, as has been mentioned, this would potentially increase their risk of sustaining injuries. In addition, players skilled in receiving the ball, passing, shooting, and decision making with the ball at their feet have more ball possession and, consequently, are exposed to more tackles and other contact situations.¹¹⁷ Furthermore, highly skilled young players are often required to compete on teams of older players. This scenario not only forces younger players to compete against more mature and physically bigger players but also to potentially play 2 matches within a very short time interval (usually less than 36 h), which may overload their immature musculoskeletal system and significantly increase their risk of injury.¹¹⁸ In this regard, Dupont et al.¹¹⁹ found that decreased recovery

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

1218

1219

1220

1221

1222

1223

1224

1225

1226

1227

1228

1229

1230

1231

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

<u>ARTICLE IN PRESS</u>

12

1245

1246

1247

1248

1249

1250

1251

1252

1253

1254

1255

1256

1257

1258

1259

1260

1261

1262

1263

1264

1265

1266

1267

1268

1269

1270

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

1281

1282

1283

1284

1285

1286

1287

1288

1289

1290

1291

1292

1293

1294

1295

1296

1297

1298

1299

1300

1301

time between matches leads to an increase in IIRs. Finally, the professionalisation of youth football has meant that many youngsters in professional academies become single-sport specialists.²⁸ High weekly training volumes associated with early specialisation may promote limited participation in other sports, decreasing motor skill development and increasing injury risk as players transition to new development cycles.^{120,121} Elite young football players who strive to be professional players may also be exposed to high levels of pressure.

However, in our study, no differences in training IIRs were found regarding the level of play for males. It is reasonable to suggest that elite players have access to better resources than do their sub-elite peers, including better equipment, comprehensive medical support, and expert coaches who can control match and training loads. These superior resources may contribute to a reduction in injury risk despite their expected greater exposure to training.¹⁷

Although elite female youth footballers showed IIRs that were similar to those shown for males, there was a lack of data related to training and matches for sub-elite female players. Future studies should analyse the injury profile for this cohort of female football players, reporting the number of injuries sustained in matches and training sessions separately.

4.7. Level and quality of the evidence

In our systematic review and meta-analysis, the pooled results of more than 25 epidemiological studies provided a moderate quality of evidence for support of the overall, training and match IIRs estimated for male youth football players. The quality of evidence for overall, training and match IIRs in females was low, coming from only 5 (training)^{27,43,49,51,53} to 10 (match)^{19,27,43,44,49,51,53,72,74,80} studies. Furthermore, several of these studies were carried out with female players who were selected to participate in various tournaments, ^{19,44,51,72,74,80} a situation that represents a shorter period of time for data collection compared to an entire football season. Therefore, future research should focus on monitoring IIRs among female youth football players throughout competitive seasons in order to provide a broader comparison with the IIRs documented among male youth footballers.

4.8. Limitations

Although our study was conducted following the international guidelines for systematic reviews and meta-analyses, some limitations should be acknowledged. Variations in injury definitions and data collection procedures used in the various studies might partly explain the heterogeneous estimates obtained in our meta-analysis and in previous meta-analyses conducted in the field of sport medicine.^{28,60,69,84} To mitigate this problem, we included in our sub-analysis only those studies that rigorously and clearly followed the time-loss injury definition described by Fuller et al.⁴¹ and Hägglund et al.⁴² The inclusion of injuries requiring medical attention may well have led to higher IIRs. However, it could also intensify the differences between data-collection procedures because non-

F.J. Robles-Palazón et al.

1302

1303

1304

1305

1306

1307

1308

1309

1310

1311

1312

1313

1314

1315

1316

1317

1318

1319

1320

1321

1322

1323

1324

time-loss IIRs have been shown to be especially sensitive to differing recording settings, and a research-invested clinical recorder might report almost a 9 times greater IIR compared to other noninvolved recorders (i.e., noninvolved physiotherapists).¹²² Thus, and based on the reality of injury surveillance among youth football players, where coaches are frequently the responsible person for recording injuries^{35,43,44,73,78} due to the lack of medical staff, a time-loss definition was used. Furthermore, because different epidemiological data were presented in the included studies (e.g., total number of injuries, number of matches played), we applied standardised formulas to account for this discrepancy. Nevertheless, even when these inclusion criteria and standardised formulas were applied, the degree of inconsistency of the main results (overall, training and match IIRs) across studies was still very high. Consequently, other aspects of football-such as differences in geographic areas or time of year affecting climatic conditions for football practice,¹²³ the monitoring period of the season,^{33,34} the number of exposure hours and match congestion,¹⁷ or the skill level of youth footballers¹¹⁷—may have constituted other sources of inconsistency. The limited number of studies reporting the location and type of injuries for elite and sub-elite players by sex made further sub-analyses in this area impossible. Such sub-analyses may have identified potential differences attributable to the level of play. However, given the results of previous studies on elite^{9,17,18,31,50,82} and sub-elite^{21,25,32-34} male and elite²⁷ and sub-elite³⁵ female players, large differences in these injury patterns might not be expected. Finally, the sample sizes of the included studies were not sufficient to investigate the interactive effects of physical maturation, growth spurt, or growth-related injuries on IIRs among young football players.

5. Conclusions

The high IIRs and probability scores found for youth footballers in our meta-analysis reinforce the need for implementing targeted injury risk-mitigation strategies in youth football, irrespective of sex. Because IIRs are higher during match play for both sexes, it is important that the training prescription mimic the demands of match play as closely as possible in order to provide the robustness and readiness needed for competitive play. The sex differences identified for the most common locations and types of injury reinforce the need for different targeted management strategies in male and female youth players. Males tend to sustain predominantly muscle injuries to the thigh, and females sustain predominantly joint and ligament injuries to the knee and ankle, so strategies should focus on neuromuscular conditioning in male players and movement mechanics, core strength, and joint stability in female players. However, there is still a paucity of data concerning female players, especially for younger and less mature female players. Additional longitudinal studies are needed to fully explore the age- and maturation-related changes in incidence, severity, location, and type of injuries that occur among footballers of both sexes.

Authors' contributions

1359

1360

1361

1362

1363

1364

1365

1366

1367

1368

1369

1370

1371

1372

1373

1374

1375

1376

1377

1378

1379

1380

1381

1382

1383

1384

1385

1386

1387

1388

1389

1390

1391

1392

1393

1394

1395

1396

1397

1398

1399

1400

1401

1402

1403

1404

1405

1406

1407

1408

1409

1410

1411

1412

1413

1414

1415

02

FJRP, ALV and FA designed the research, conducted the searches and screening process, extracted the data, interpreted the data analysis, and led the drafting of the manuscript; AGG performed all the statistical analyses; MDSC, JLO and PSdB assisted in the interpretation of the data analysis and contributed to the drafting of the manuscript. All authors have read and approved the final version of the manuscript and agree with the order of presentation of the authors.

Competing interests

The authors declare that they have no competing interests.

Acknowledgments

The authors thank all the researchers who provided additional information. Part of this work was carried out during a research stay of the first author at Cardiff Metropolitan University from March 2020 to July 2020. Francisco Javier Robles-Palazón was supported by the Program of Human Resources Formation for Science and Technology (20326/FPI/2017) from the Seneca Foundation-Agency for Science and Technology in the Region of Murcia (Spain). Francisco Ayala was supported by a Ramón y Cajal postdoctoral fellowship given by the Spanish Ministry of Science and Innovation (RYC2019-028383-I). This study is part of the project titled Estudio del riesgo de lesión en jóvenes deportistas a través de redes de inteligencia artificial, funded by the Spanish Ministry of Science and Innovation (DEP2017-88775-P), the State Research Agency (AEI), and the European Regional Development Fund (ERDF). The funders had no role in the study's design, data analysis, interpretation, or decision to submit the work for publication.

Supplementary materials

Supplementary material associated with this article can be found in the online version at doi:10.1016/j.jshs.2021.10.002.

References

- Hulteen RM, Smith JJ, Morgan PJ. Global participation in sport and leisure-time physical activities: a systematic review and meta-analysis. *Prev Med* 2017;95:14–25.
- Krustrup P, Aagaard P, Nybo L, Petersen J, Mohr M, Bangsbo J. Recreational football as a health promoting activity: a topical review. *Scand J Med Sci Sports* 2010;20:1–13.
- Faude O, Rössler R, Junge A. Football injuries in children and adolescent players: are there clues for prevention? *Sports Med* 2013;43:819–37.
- Hjelm N, Werner S, Renstrom P. Injury profile in junior tennis players: a prospective two year study. *Knee Surg Sports Traumatol Arthrosc* 2010;18:845–50.
- Caine D, Knutzen K, Howe W, et al. A three-year epidemiological study of injuries affecting young female gymnasts. *Phys Ther Sport* 2003;4:10–23.
- Caine D, Maffulli N, Caine C. Epidemiology of injury in child and adolescent sports: injury rates, risk factors, and prevention. *Clin Sports Med* 2008;27:19–50.

- 7. Emery CA, Meeuwisse WH, McAllister JR. Survey of sport participation and sport injury in Calgary and area high schools. *Clin J Sport Med* 2006;**16**:20–6.
- Read PJ, Oliver JL, De Ste Croix MBA, Myer GD, Lloyd RS. An audit of injuries in six english professional soccer academies. *J Sports Sci* 2018;36:1542–8.
- Renshaw A, Goodwin PC. Injury incidence in a Premier League youth soccer academy using the consensus statement: a prospective cohort study. *BMJ Open Sport Exerc Med* 2016;2: e000132. doi:10.1136/ bmjsem-2016-000132.
- van der Sluis A, Elferink-Gemser MT, Coelho-e-Silva MJ, Nijboer JA, Brink MS, Visscher C. Sport injuries aligned to peak height velocity in talented pubertal soccer players. *Int J Sports Med* 2014;35:351–5.
- Kolstrup LA, Koopmann KU, Nygaard UH, Nygaard RH, Agger P. Injuries during football tournaments in 45,000 children and adolescents. *Eur J Sport Sci* 2016;16:1167–75.
- Maffulli N, Longo UG, Gougoulias N, Loppini M, Denaro V. Long-term health outcomes of youth sports injuries. *Br J Sports Med* 2010;44:21–5.
- Myer GD, Faigenbaum AD, Ford KR, Best TM, Bergeron MF, Hewett TE. When to initiate integrative neuromuscular training to reduce sportsrelated injuries and enhance health in youth? *Curr Sports Med Rep* 2011;**10**:155–66.
- Koutures CG, Gregory AJM. Clinical report: injuries in youth soccer. *Pediatrics* 2010;**125**:410–4.
- van Mechelen W, Hlobil H, Kemper HC. Incidence, severity, aetiology and prevention of sports injuries. Sports Med 1992;14:82–99.
- Finch C. A new framework for research leading to sports injury prevention. J Sci Med Sport 2006;9:3–9.
- 17. Tears C, Chesterton P, Wijnbergen M. The elite player performance plan: the impact of a new national youth development strategy on injury characteristics in a premier league football academy. *J Sports Sci* 2018;**36**:2181–8.
- Raya-González J, Suárez-Arrones L, Navandar A, Balsalobre-Fernández C, Sáez de Villarreal E. Injury profile of elite male young soccer players in a Spanish male professional soccer club: a prospective study during four consecutive seasons. *J Sport Rehabil* 2019;29:801–7.
- Lislevand M, Andersen TE, Junge A, Dvorak J, Steffen K. Injury surveillance during a 2-day national female youth football tournament in Kenya. *Br J Sports Med* 2014;48:924–8.
- Nilsson T, Östenberg AH, Alricsson M. Injury profile among elite male youth soccer players in a Swedish first league. J Exerc Rehabil 2016;12:83–9.
- Chena-Sinovas M, ML Rodríguez-Hernández. Bores-Cerezal A. Epidemiology of injuries in young Spanish soccer players. *Med dello Sport* 2019;72:254–66.
- Cezarino LG, da Silva Grüninger BL, Silva RS. Injury profile in a Brazilian first-division youth soccer team: a prospective study. *J Athl Train* 2020;55:295–302.
- Bult HJ, Barendrecht M, Tak IJR. Injury risk and injury burden are related to age group and peak height velocity among talented male youth soccer players. *Orthop J Sports Med* 2018;6: 2325967118811042. doi:10.1177/2325967118811042.
- Bianco A, Spedicato M, Petrucci M, et al. A prospective analysis of the injury incidence of young male professional football players on artificial turf. *Asian J Sports Med* 2016;7:e28425. doi:10.5812/asjsm.28425.
- Nogueira M, Laiginhas R, Ramos J, Costa O. Injuries in Portuguese amateur youth football players: a six month prospective descriptive study. *Acta Med Port* 2017;30:840–7.
- Clausen MB, Zebis MK, Møller M, et al. High injury incidence in adolescent female soccer. *Am J Sports Med* 2014;42:2487–94.
- 27. Le Gall F, Carling C, Reilly T. Injuries in young elite female soccer players: an 8-season prospective study. *Am J Sports Med* 2008;**36**:276–84.
- Jones S, Almousa S, Gibb A, et al. Injury incidence, prevalence and severity in high-level male youth football: a systematic review. *Sports Med* 2019;49:1879–99.
- 29. Hewett TE, Myer GD, Ford KR. Anterior cruciate ligament injuries in female athletes: part 1, mechanisms and risk factors. *Am J Sports Med* 2006;**34**:299–311.

1416

1417

1418

1419

1420

1421

1422

1423

1424

1425

1426

1427

F.J. Robles-Palazón et al.

1530

1531

1532

1533

1534

1535

1536

1537

1538

1539

1540

1541

1542

1543

1544

1545

1546

1547

1548

1549

1550

1551

1552

1553

1554

1555

1556

1557

1558

1559

1560

1561

1562

1563

1564

1565

1566

1567

1568

1569

1570

1571

1572

1573

1574

1575

1576

1577

1578

1579

1580

1581

1582

1583

1584

1585

1586

14

Athl Train 2004;39:352-64.

2013;47:339-46.

2010:44:809-15.

cochrane.org.

31. Ergün M, Denerel HN, Binnet MS, Ertat KA. Injuries in elite youth foot-

32. Frisch A, Urhausen A, Seil R, Croisier JL, Windal T, Theisen D. Associ-

33. Brito J, Malina RM, Seabra A, et al. Injuries in Portuguese youth soccer

players during training and match play. J Athl Train 2012;47:191-7.

34. Brito J, Rebelo A, Soares JM, Seabra A, Krustrup P, Malina RM. Injuries

35. Soderman K, Adolphson J, Lorentzon R, Alfredson H. Injuries in adoles-

36. Watson A, Brickson S, Brooks A, Dunn W. Subjective well-being and

37. Brink MS, Visscher C, Arends S, Zwerver J, Post WJ, Lemmink KA.

38. Price RJ, Hawkins RD, Hulse MA, Hodson A. The Football Association

39. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items

40. Higgins JPT, Green S. Cochrane Handbook for Systematic Reviews of

41. Fuller CW, Ekstrand J, Junge A, et al. Consensus statement on injury

42. Hägglund M, Waldén M, Bahr R, Ekstrand J. Methods for epidemiologi-

43. Kuzuhara K, Shibata M, Uchida R. Injuries in Japanese junior soccer

44. Soligard T, Bahr R, Andersen TE. Injury risk on artificial turf and grass

45. Sieland J, Krause F, Kalo K, et al. Injuries and functional performance

46. Fouasson-Chailloux A, Mesland O, Menu P, Dauty M. Soccer injuries

47. Azuma N, Someya F. Injury prevention effects of stretching exercise

48. Rommers N, Rössler R, Verhagen E, et al. A machine learning approach

49. Sprouse B, Alty J, Kemp S, et al. The football association injury and ill-

50. Wik EH, Lolli L, Chamari K, et al. Injury patterns differ with age in male

51. Hägglund M, Waldén M, Ekstrand J. UEFA injury study - An injury

52. Brito JJ, Rebelo AA, Soares JMJM, Seabra AA, Krustrup P, Malina RM.

in an elite national academy. Br J Sports Med 2020;55:794-800.

players: a 5-year cohort study. Sci Sports 2020;35:145-53.

in youth tournament football. Scand J Med Sci Sports 2012;22:356-61.

status in young elite football players: a prospective 2-year monitoring. J

documented by F-MARC guidelines in 13- and 14-year old national elite

intervention by physical therapists in male high school soccer players.

to assess injury risk in elite youth football players. Med Sci Sports Exerc

ness surveillance study: the incidence, burden and severity of injuries

and illness in men's and women's international football. Sports Med.

vouth football: a four-season prospective study of 1111 time-loss injuries

audit of European Championships 2006 to 2008. Br J Sports Med

Injuries in youth soccer during the preseason. Clin J Sport Med

players during games and practices. J Athl Train 2017;52:1147-52.

outdoor soccer season. Scand J Med Sci Sports 2001;11:299-304.

soccer players. Br J Sports Med 2017;51:194-9.

ball. Br J Sports Med 2004;38:466-71.

injuries. Br J Sports Med 2006;40:193-201.

UEFA model. Br J Sports Med 2005;39:340-6.

Sports Med Phys Fitness 2020;60:1363-70.

Scand J Med Sci Sports 2020;30:2178-92.

2020. doi:10.1007/s40279-020-01411-8.

2020;52:1745-51.

2009;43:483-9.

2011;21:259-60.

Intern Med 2009;151:264-9.

in youth soccer during the preseason. Clin J Sport Med 2011;21:259-60.

cent female players in European football: a prospective study over one

training load predict in-season injury and illness risk in female youth

Monitoring stress and recovery: new insights for the prevention of inju-

ries and illnesses in elite youth soccer players. Br J Sports Med

medical research programme: an audit of injuries in academy youth foot-

for systematic reviews and meta-analyses: The PRISMA statement. Ann

Interventions Version 5.1.0 [Updated March 2011]. London, UK: The

Cochrane Collaboration; 2011.. Available from; . www.handbook.

definitions and data collection procedures in studies of football (soccer)

cal study of injuries to professional football players: developing the

prospective follow-up. Scand J Med Sci Sports 2011;21:e468-76.

ball players: a prospective three-year study. Acta Orthop Traumatol Turc

ation between preseason functional tests and injuries in youth football: a

- 1489 1490
- 1491 1492
- 1493
- 1494
- 1495 1496
- 1497 1498
- 1499 1500

1501

1502 1503

1504 1505

1506 1507

1508 1509

1510

1511 1512 1513

1514

1515 1516

1517

1518

1519 1520 1521

1522 1523 1524

1525

1526

1527 1528 1529

- 30. Myer GD, Ford KR, Hewett TE. Rationale and clinical techniques for 53. Soligard T, Myklebust G, Steffen K, et al. Comprehensive warm-up proanterior cruciate ligament injury prevention among female athletes. J gramme to prevent injuries in young female footballers: cluster randomised controlled trial. BMJ 2008;337:1-9.
 - 54. Imai A, Imai T, Iizuka S, Kaneoka K. A trunk stabilization exercise warm-up may reduce ankle injuries in junior soccer players. Int J Sports Med 2018;39:270-4.
 - 55. Delecroix B, Delaval B, Dawson B, Berthoin S, Dupont G. Workload and injury incidence in elite football academy players. J Sports Sci 2019:37:2768-73.
 - 56. Hawkins RD, Fuller CW. A prospective epidemiological study of injuries in four English professional football clubs. Br J Sports Med 1999:33:196-203.
 - 57. Lefebvre C, Glanville J, Briscoe S, et al. Technical Supplement to Chapter 4: searching for and selecting studies. In: Higgins JPT, Thomas J, Chandler J, editors. Cochrane Handbook for Systematic Reviews of Interventions. London, UK: The Cochrane Collaboration; 2019.p.1-94.
 - 58. Pfirrmann D, Herbst M, Ingelfinger P, Simon P, Tug S. Analysis of injury incidences in male professional adult and elite youth soccer players: a systematic review. J Athl Train 2016;51:410-24.
 - 59. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. Int J Surg 2014;12:1495-9.
 - 60. López-Valenciano A, Ruiz-Pérez I, Garcia-Gómez A, et al. Epidemiology of injuries in professional football: a systematic review and metaanalysis. Br J Sports Med 2020;54:711-8.
 - 61. Orwin RG, Vevea JL. Evaluating coding decisions. In: Cooper H, Hedges LV, Valentine JC, editors. The Handbook of Research Synthesis and Meta-Analysis. New York, NY: Russell Sage Foundation; 2009. p.177-203
 - 62. Guyatt GH, Oxman AD, Montori V, et al. GRADE guidelines: 5. Rating the quality of evidence-publication bias. J Clin Epidemiol 2011;64:1277-82.
 - 63. Balshem H, Helfand M, Schünemann HJ, et al. GRADE guidelines: 3. Rating the quality of evidence. J Clin Epidemiol 2011;64:401-6.
 - 64. Williams S, Trewartha G, Kemp S, Stokes K. A meta-analysis of injuries in senior men's professional Rugby Union. Sports Med 2013;43:1043-55.
 - 65. Lystad RP, Pollard H, Graham PL. Epidemiology of injuries in competition taekwondo: a meta-analysis of observational studies. J Sci Med Sport 2009;12:614-21.
 - 66. Bagos PG, Nikolopoulos GK. Mixed-effects Poisson regression models for meta-analysis of follow-up studies with constant or varying durations. Int J Biostat 2009;5:21. doi:10.2202/1557-4679.1168.
 - 67. Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. Stat Med 2002;21:1539-58.
 - 68. Parekh N, Hodges SD, Pollock AM, Kirkwood G. Communicating the risk of injury in schoolboy rugby: using Poisson probability as an alternative presentation of the epidemiology. Br J Sports Med 2012;46:611-3.
 - 69. Freitag A, Kirkwood G, Scharer S, Ofori-Asenso R, Pollock AM. Systematic review of rugby injuries in children and adolescents under 21 years. Br J Sports Med 2015;49:511-9.
 - 70. Nevill AM, Atkinson G, Hughes MD, Cooper SM. Statistical methods for analysing discrete and categorical data recorded in performance analysis. J Sports Sci 2002;20:829-44.
 - 71. Viechtbauer W. Conducting meta-analyses in R with the metafor package. J Stat Softw 2010;36:i03. doi:10.18637/jss.v036.i03.
 - 72. Andreasen I, Fauno P, Lund B, Lemche P, Knudsen H. Soccer injuries among youth. Scand J Med Sci Sports 1993;3:62-6.
 - 73. Backous DD, Friedl KE, Smith NJ, Parr TJ, Carpine WD. Soccer injuries and their relation to physical maturity. Am J Dis Child 1988;142:839-42.
 - Junge A, Dvorak J. Injury surveillance in the world football tournaments 74. 1998-2012. Br J Sports Med 2013;47:782-8.
 - 75. Junge A, Cheung K, Edwards T, Dvorak J. Injuries in youth amateur soccer and rugby players-comparison of incidence and characteristics. Br J Sports Med 2004;38:168-72.
 - 76. Junge A, Rösch D, Peterson L, Graf-Baumann T, Dvorak J. Prevention of soccer injuries: a prospective intervention study in youth amateur players. Am J Sports Med 2002;30:652-9.

Please cite this article as: Francisco Javier Robles-Palazón et al., Epidemiology of injuries in male and female youth football players: A systematic review and meta-analysis, Journal of Sport and Health Science (2021), https://doi.org/10.1016/j.jshs.2021.10.002

Epidemiology of injuries in youth football

1587

1588

1589

1590

1591

1592

1593

1594

1595

1596

1597

1598

1599

1600

1601

1602

1603

1604

1605

1606

1607

1608

1609

1610

1611

1612

1613

1614

1615

1616

1617

1618

1619

1620

1621

1622

1623

1624

1625

1626

1627

1628

1629

1630

1631

1632

1633

1634

1635

1636

1637

1638

1639

1640

1641

1642

1643

- Kakavelakis KN, Vlazakis S, Vlahakis I, Charissis G. Soccer injuries in childhood. Scand J Med Sci Sports 2003;13:175–8.
- Nielsen AB, Yde J. Epidemiology and traumatology of injuries in soccer. *Am J Sports Med* 1989;17:803–7.
- Owoeye OBA, Akinbo SRA, Tella BA, Olawale OA. Efficacy of the FIFA 11+ warm-up programme in male youth football: a cluster randomised controlled trial. J Sport Sci Med 2014;13:321–8.
- Schmidt-Olsen S, Bünemann LKH, Lade V, Brassoe JOK. Soccer injuries of youth. *Br J Sports Med* 1985;19:161–4.
- Steffen K, Myklebust G, Olsen OE, Holme I, Bahr R. Preventing injuries in female youth football: a cluster-randomized controlled trial. *Scand J Med Sci Sports* 2008;18:605–14.
- Waldén M, Hägglund M, Ekstrand J. Football injuries during European Championships 2004–2005. *Knee Surg Sports Traumatol Arthrosc* 2007;15:1155–62.
- 83. Zarei M, Abbasi H, Namazi P, Asgari M, Rommers N, Rössler R. The 11 + Kids warm-up programme to prevent injuries in young Iranian male high-level football (soccer) players: a cluster-randomised controlled trial. J Sci Med Sport 2020;23:469–74.
- López-Valenciano A, Raya-González J, García-Gómez JA, et al. Injury profile in women's football: a systematic review and meta-analysis. *Sports Med* 2021;51:423–42.
- Olsen OE, Myklebust G, Engebretsen L, Bahr R. Injury pattern in youth team handball: a comparison of two prospective registration methods. *Scand J Med Sci Sports* 2006;16:426–32.
- Longo UG, Loppini M, Berton A, Marinozzi A, Maffulli N, Denaro V. The FIFA 11+ program is effective in preventing injuries in elite male basketball players: a cluster randomized controlled trial. *Am J Sports Med* 2012;40:996–1005.
- Bere T, Kruczynski J, Veintimilla N, Hamu Y, Bahr R. Injury risk is low among world-class volleyball players: 4-year data from the FIVB Injury Surveillance System. *Br J Sports Med* 2015;49:1132–7.
- Bangsbo J, Mohr M, Krustrup P. Physical and metabolic demands of training and match-play in the elite football player. *J Sports Sci* 2006;24:665–74.
- Giannotti M, Al-Sahab B, McFaull S, Tamim H. Epidemiology of acute soccer injuries in Canadian children and youth. *Pediatr Emerg Care* 2011;27:81–5.
- Blackburn JT, Padua DA. Influence of trunk flexion on hip and knee joint kinematics during a controlled drop landing. *Clin Biomech* 2008;23:313–9.
- Raya-González J, Suarez-Arrones L, Larruskain J, Sáez de Villarreal E. Muscle injuries in the academy of a Spanish professional football club: a one-year prospective study. *Apunt Med l'Esport* 2018;53:3–9.
- Martínez-Lagunas V, Niessen M, Hartmann U. Women's football: player characteristics and demands of the game. J Sport Health Sci 2014;3:258–72.
- Beaudouin F, Gioftsidou A, Larsen MN, et al. The UEFA Heading Study: heading incidence in children's and youth football (soccer) in eight European countries. *Scand J Med Sci Sports* 2020;**30**:1506–17.
- Faude O, Rössler R, Junge A, et al. Head injuries in children's footballresults from two prospective cohort studies in four European countries. *Scand J Med Sci Sports* 2017;27:1986–92.
- Mooney J, Self M, Refaey K, et al. Concussion in soccer: a comprehensive review of the literature. *Concussion* 2020;5:CNC76. doi:10.2217/ cnc-2020-0004.
- 96. Rössler R, Junge A, Bizzini M, et al. A multinational cluster randomised controlled trial to assess the efficacy of "11 + Kids": a warm-up programme to prevent injuries in children's football. *Sports Med* 2018;48:1493–504.
- Steib S, Rahlf AL, Pfeifer K, Zech A. Dose-response relationship of neuromuscular training for injury prevention in youth athletes: a meta-analysis. *Front Physiol* 2017;8:920. doi:10.3389/fphys.2017.00920.
- Eirale C, Tol JL, Farooq A, Smiley F, Chalabi H. Low injury rate strongly correlates with team success in Qatari professional football. Br J Sports Med 2013;47:807–8.
- Le Gall F, Carling C, Reilly T. Biological maturity and injury in elite youth football. Scand J Med Sci Sports 2007;17:564–72.
- 100. Materne O, Chamari K, Farooq A, et al. Injury incidence and burden in a youth elite football academy: a four-season prospective study of 551 players aged from under 9 to under 19 years. *Br J Sports Med* 2021;55:493–500.

- 101. Le Gall F, Carling C, Reilly T, Vandewalle H, Church J. Rochcongar P. Incidence of injuries in elite French youth soccer players: a 10-season study. *Am J Sports Med* 2006;**34**:928–38.
- Emery CA, Meeuwisse WH, Hartmann SE. Evaluation of risk factors for injury in adolescent soccer: implementation and validation of an injury surveillance system. *Am J Sports Med* 2005;**33**:1882–91.
- 103. Kucera KL, Marshall SW, Kirkendall DT, Marchak PM, Garrett Jr WE. Injury history as a risk factor for incident injury in youth soccer. Br J Sports Med 2005;39:462–6.
- 104. Wiggins AJ, Grandhi RK, Schneider DK, Stanfield D, Webster KE, Myer GD. Risk of secondary injury in younger athletes after anterior cruciate ligament reconstruction: a systematic review and meta-analysis. *Am J Sports Med* 2016;44:1861–76.
- López-Valenciano A, Ayala F, Puerta JM, et al. A preventive model for muscle injuries: a novel approach based on learning algorithms. *Med Sci Sports Exerc* 2018;**50**:915–27.
- 106. Hägglund M, Waldén M, Ekstrand J. Previous injury as a risk factor for injury in elite football: a prospective study over two consecutive seasons. *Br J Sports Med* 2006;**40**:767–72.
- 107. Faude O, Rössler R, Petushek EJ, Roth R, Zahner L, Donath L. Neuromuscular adaptations to multimodal injury prevention programs in youth sports: a systematic review with meta-analysis of randomized controlled trials. *Front Physiol* 2017;8:791. doi:10.3389/fphys.2017.00791.
- Gabbe BJ, Bailey M, Cook JL, et al. The association between hip and groin injuries in the elite junior football years and injuries sustained during elite senior competition. *Br J Sports Med* 2010;44:799–802.
- 109. Moran J, Sandercock GRH, Ramírez-Campillo R, Meylan C, Collison J, Parry DA. A meta-analysis of maturation-related variation in adolescent boy athletes' adaptations to short-term resistance training. *J Sports Sci* 2017;**35**:1041–51.
- 110. Faigenbaum AD, Lloyd RS, Oliver JL. *Essentials of Youth Fitness*. Champaign, IL: Human Kinetics Publishers; 2019.
- 111. Ratel S, Duché P, Williams CA. Muscle fatigue during high-intensity exercise in children. *Sports Med* 2006;**36**:1031–65.
- 112. Philippaerts RM, Vaeyens R, Janssens M, et al. The relationship between peak height velocity and physical performance in youth soccer players. J Sports Sci 2006;24:221–30.
- 113. Kemper GLJ, van der Sluis A, Brink MS, Visscher C, Frencken WGP, Elferink-Gemser MT. Anthropometric injury risk factors in elite-standard youth soccer. *Int J Sports Med* 2015;36:1112–7.
- 114. Buchheit M, Horobeanu C, Mendez-Villanueva A, Simpson BM, Bourdon PC. Effects of age and spa treatment on match running performance over two consecutive games in highly trained young soccer players. J Sports Sci 2011;29:591–8.
- 115. Buchheit M, Mendez-Villanueva A, Simpson BM, Bourdon PC. Match running performance and fitness in youth soccer. *Int J Sports Med* 2010;**31**:818–25.
- Bowen L, Gross AS, Gimpel M, Li FX. Accumulated workloads and the acute: chronic workload ratio relate to injury risk in elite youth football players. *Br J Sports Med* 2017;**51**:452–9.
- 117. Soligard T, Grindem H, Bahr R, Andersen TE. Are skilled players at greater risk of injury in female youth football? *Br J Sports Med* 2010;**44**:1118–23.
- Nilstad A, Andersen TE, Bahr R, Holme I, Steffen K. Risk factors for lower extremity injuries in elite female soccer players. *Am J Sports Med* 2014;42:940–8.
- Dupont G, Nedelec M, McCall A, McCormack D, Berthoin S, Wisloff U. Effect of 2 soccer matches in a week on physical performance and injury rate. *Am J Sports Med* 2010;**38**:1752–8.
- 120. Jayanthi NA, LaBella CR, Fischer D, Pasulka J, Dugas LR. Sports-specialized intensive training and the risk of injury in young athletes: a clinical case-control study. *Am J Sports Med* 2015;**43**:794–801.
- 121. Mostafavifar AM, Best TM, Myer GD. Early sport specialisation, does it lead to long-term problems? *Br J Sports Med* 2013;**47**:1060–1.
- 122. Wik EH, Materne O, Chamari K, et al. Involving research-invested clinicians in data collection affects injury incidence in youth football. *Scand J Med Sci Sports* 2019;29:1031–9.
- 123. Watson A, Mjaanes JM. Soccer injuries in children and adolescents. *Pediatrics* 2019;144: e20192759. doi:10.1542/peds.2019-2759.

1644

1680

1681

1682

1683

1684

1685

1686

1687

1688

1689

1690

1691

1692

1693

1694

1695

1696

1697

1698

1699